

Package: SorptionAnalysis (via r-universe)

August 20, 2024

Type Package

Title Static Adsorption Experiment Plotting and Analysis

Version 0.1.0

Author Aurnov Chattopadhyay

Maintainer Aurnov Chattopadhyay <aurnovcy@gmail.com>

Description Provides tools to efficiently analyze and visualize laboratory data from aqueous static adsorption experiments. The package provides functions to plot Langmuir, Freundlich, and Temkin isotherms and functions to determine the statistical conformity of data points to the Langmuir, Freundlich, and Temkin adsorption models through statistical characterization of the isothermic least squares regressions lines. Scientific Reference: Dada, A.O, Olalekan, A., Olatunya, A. (2012) <doi:10.9790/5736-0313845>.

License GPL-2

Encoding UTF-8

LazyData true

RoxygenNote 6.0.1

NeedsCompilation no

Repository <https://aurnovcy.r-universe.dev>

RemoteUrl <https://github.com/aurnovcy/sorptionanalysis>

RemoteRef HEAD

RemoteSha 386f871cab7e7c57a61ca9fb8872e05b80b2e510

Contents

freundlichanalysis	2
freundlichplot	2
langmuiranalysis	3
langmuirplot	4
summaryanalysis	4

summaryplots	5
temkinanalysis	5
temkinplot	6

Index	7
--------------	----------

freundlichanalysis	<i>Describes the conformity of the results from a static adsorption experiment to the Freundlich Isotherm model</i>
--------------------	---

Description

Describes the conformity of the results from a static adsorption experiment to the Freundlich Isotherm model

Usage

freundlichanalysis(Ce, Qe)

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The residuals, y - intercept, coefficient, and correlation factor of the LSRL based on the Temkin isotherm model

Examples

freundlichanalysis(c(1,2,3,4), c(1,2,3,4))

freundlichplot	<i>Plots a Freundlich Isotherm using results from a static adsorption experiment</i>
----------------	--

Description

Plots a Freundlich Isotherm using results from a static adsorption experiment

Usage

freundlichplot(Ce, Qe)

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The plot of $\log(C_e)$ against $\log(Q_e)$ with the LSRL displayed

Examples

```
freundlichplot(c(1,2,3,4), c(1,2,3,4))
```

langmuiranalysis	<i>Describes the conformity of the results from a static adsorption experiment to the Langmuir Isotherm model</i>
------------------	---

Description

Describes the conformity of the results from a static adsorption experiment to the Langmuir Isotherm model

Usage

```
langmuiranalysis(Ce, Qe)
```

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The residuals, y - intercept, coefficient, and correlation factor of the LSRL based on the Temkin isotherm model

Examples

```
langmuiranalysis(c(1,2,3,4), c(1,2,3,4))
```

langmuirplot	<i>Plots a Langmuir Isotherm using results from a static adsorption experiment</i>
--------------	--

Description

Plots a Langmuir Isotherm using results from a static adsorption experiment

Usage

```
langmuirplot(Ce, Qe)
```

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The plot of 1 over Ce against 1 over Qe with the LSRL displayed

Examples

```
langmuirplot(c(1,2,3,4), c(1,2,3,4))
```

summaryanalysis	<i>Describes the conformity of the results from a static adsorption experiment to the Langmuir, Freundlich, and Temkin Isotherm model</i>
-----------------	---

Description

Describes the conformity of the results from a static adsorption experiment to the Langmuir, Freundlich, and Temkin Isotherm model

Usage

```
summaryanalysis(Ce, Qe)
```

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The residuals, y - intercept, coefficient, and correlation factor of the LSRL based on the Langmuir, Freundlich, and Temkin isotherm model

Examples

```
summaryanalysis(c(1,2,3,4), c(1,2,3,4))
```

summaryplots	<i>Plots the three different Isotherm models using results from a static adsorption experiment</i>
--------------	--

Description

Plots the three different Isotherm models using results from a static adsorption experiment

Usage

```
summaryplots(Ce, Qe)
```

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The plots of Langmuir, Freundlich, and Temkin Isotherms

Examples

```
summaryplots(c(1,2,3,4), c(1,2,3,4))
```

temkinanalysis	<i>Describes the conformity of the results from a static adsorption experiment to the Temkin Isotherm model</i>
----------------	---

Description

Describes the conformity of the results from a static adsorption experiment to the Temkin Isotherm model

Usage

```
temkinanalysis(Ce, Qe)
```

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The residuals, y - intercept, coefficient, and correlation factor of the LSRL based on the Temkin isotherm model

Examples

```
temkinanalysis(c(1,2,3,4), c(1,2,3,4))
```

temkinplot	<i>Plots a Temkin Isotherm using results from a static adsorption experiment</i>
------------	--

Description

Plots a Temkin Isotherm using results from a static adsorption experiment

Usage

```
temkinplot(Ce, Qe)
```

Arguments

Ce	A numeric vector consisting of equilibrium concentration values
Qe	A numeric vector consisting of quantities adsorbed at equilibrium

Value

The plot of $\ln(Ce)$ against Qe with the LSRL displayed

Examples

```
temkinplot(c(1,2,3,4), c(1,2,3,4))
```

Index

freundlichenanalysis, [2](#)
freundlichplot, [2](#)

langmuiranalysis, [3](#)
langmuirplot, [4](#)

summaryanalysis, [4](#)
summaryplots, [5](#)

temkinanalysis, [5](#)
temkinplot, [6](#)